

08/699716



08/27/96

08/699716

TRANSMITTAL LETTER

To the Commissioner of Patents and Trademarks:

003/029/SAP
Client/MatterTransmitted herewith for filing is the X Utility Design Plant patent application of:Inventor(s) 1996 G. Smith, Arthur M. Friedlander, George W. Anderson, Susan L. WelkosTitle of invention Recombinant F1-V Plague Vaccine

Enclosed are the following items:

- X Duplicate copies of this Patent Application Transmittal form.
2 Sheet(s) of (X) informal () formal drawings.
35 Page(s) of specification.
5 Page(s) of claims.
1 Page(s) of abstract.
X Declaration (X) signed () unsigned.
 A Power of Attorney executed by the () inventors () Assignee.
 An assignment transmittal letter.
 A verified statement to establish Small Entity Status under 37 CFR 1.9 and 37 CFR 1.27.
X An information Disclosure Statement.
X A list of references (Form-1449).
X Copies of references.
 Preliminary amendment.
X Postcard.
X Other (specify) COMPUTER READABLE FORM OF SEQUENCE LISTING

CLAIMS AS FILED

BASIC FILING FEE			SMALL ENTITY	OR	OTHER THAN SMALL ENTITY	
<u>X</u> UTILITY			\$ 375	OR	\$ 750	\$ 750
<u> </u> DESIGN			\$ 155	OR	\$ 310	\$ -
<u> </u> PLANT			\$ 255	OR	\$ 510	\$ -
TOTAL CLAIMS	<u>28</u> - 20 = <u>8</u>	X	\$ 11	OR	\$ 22	= \$ 176
INDEPENDENT CLAIMS	<u>13</u> - 3 = <u>10</u>	X	\$ 39	OR	\$ 78	= \$ 780
MULTIPLE DEPENDENT CLAIMS PRESENT		+	\$ 115	OR	\$ 230	= \$ -
RECORDING FEE FOR ASSIGNMENT (\$ 40.00/EA.)						\$ -
TOTAL FILING FEE:						\$ 1,706

The Commissioner is hereby authorized to charge and credit Deposit Account No. 21-0380 as described below:

- X Charge the total filing fee of \$ 1,706.
X Charge any additional filing fees as required under 37 CFR 1.16 and 1.17.
X Credit any overpayment.
X Charge the issue fee required under 37 CFR 1.18 at the time of mailing of the Notice of Allowance, pursuant to 37 CFR 1.311(b).

Claim for Priority Under 35 U.S.C., Section 120:

() The benefit under 35 U.S.C., Section 120 is hereby claimed from the following United States application:

Claim for Priority Under 35 U.S.C., Section 119:

() The benefit of priority under 35 U.S.C. Section 119 is hereby claimed from the following foreign application:

Please address all correspondence in connection with this application to:

JOHN F. MORAN
 U. S. Army Medical Research and Materiel Command
 ATTN: MCMR-JA (JOHN MORAN)
 Fort Detrick, Frederick, Maryland 21702-5012
 Telephone: (301) 619-2065

By: John F. Moran No 39,441

John F. Moran, Reg. No. 26,313

Date: Aug 27, 1996

SAP 8/94

08699716-082796

jc965 U.S. PTO

08/699716



08/27/96

08/699716



APPLICATION FOR UNITED STATES PATENT

INVENTOR(S): David G. Heath
Arthur M. Friedlander
George W. Anderson
Susan L. Welkos

INVENTION: RECOMBINANT F1-V PLAGUE VACCINE

SPECIFICATION

SAP 9/94

08/699716-08296

jc965 U.S. PTO
08/699716
08/27/96

A. Segue
08/699716



TITLE OF THE INVENTION

Recombinant F1-V Plague Vaccine

by

David G. Heath
Arthur M. Friedlander
George W. Anderson
Susan L. Welkos

10

INTRODUCTION

15 *Yersinia pestis* is the causative organism of plague in a wide range of animals including man. Bubonic plague in man is most commonly acquired from the bite of an infected flea and is characterized by the swelling of local lymph nodes which form buboes. One of the complications of bubonic plague is secondary pneumonia, and in these cases plague can be transmitted between humans by airborne droplets. Pneumonic plague, particularly, is extremely virulent and results in high mortality rates.

20 Plague is endemic in regions of North and South America, Africa, China and Asia and, as evidenced by the recent outbreak of pneumonic plague in India [Centers for Disease Control. (1994). Update: Human Plague-India. *MMWR* 43:722-723], epidemics of enormous consequences remain a potential for this organism. Thus, there is a clear need
25 for a vaccine which would protect individuals living and traveling in endemic areas.

30 The current human, licensed, vaccines available for prevention of plague are whole cell vaccines. A number of formulations exist. The plague USP vaccine, comprising formaldehyde killed *Y. pestis* bacilli, which is administered to the body via intramuscular injection, produces local and systemic side-effects, ranging from mild headaches to severe malaise and fever. Additionally, the vaccine does not provide complete immunity, since vaccinated individuals can contract pneumonic plague, indicating inadequate immunity at mucosal surfaces.

 The live attenuated vaccine EV76 [Meyer *et al.* (1974) *J. Infect. Dis.* 129 suppl., 13-18] was tested extensively and used in the former Soviet Union from 1939, although its

08699716 "082796

efficacy in evoking an immune response in man is questionable [Meyer *et al.* (1974) *J. Infect. Dis.* **129** suppl., 85-120]. The virulence of EV76 differs in several animal species, and non-human primates are particularly susceptible to a chronic infection with this strain. In the Western World, the vaccine is considered to be unsuitable for mass vaccination due
5 to the severity of the side-effects and the possibility of the strain reverting to full virulence.

Efforts to develop a more effective vaccine have focused on acellular subunits from *Y. pestis* as immunogens. Two of the candidate subunits are the F1 and V antigens. The capsule surrounding *Y. pestis* cells is composed of a protein component known as Fraction 1 (F1) [Baker *et al.* (1952) *J. Immunol.* **68**: 131-145] which is only fully expressed at
10 37°C and encoded on the 100 kb pFra plasmid [Protsenko *et al.* (1983) *Genetika* **19**: 1081-1090]. This complex confers resistance to phagocytosis. Detection of antibodies to F1 is the basis of standard serological tests for the surveillance and diagnosis of plague as infected animals and humans produce a strong humoral response to the antigen [Shepherd
15 *et al.* (1986) *J. Clin. Microbiol.* **24**: 1075-1078; Williams *et al.* (1982) *Bull. World Health Organ.* **64**: 745-752].

V antigen, postulated to act as a virulence factor, is a 37 kDa secreted protein which acts as a cytoplasmic regulator of Yops (Yersinia outer membrane protein) expression. The V antigen is encoded on a homologous 75 kb low-calcium response (LCR) plasmid present in *Y. pestis*, *Y. pseudotuberculosis* and *Y. enterocolitica*. This plasmid mediates the
20 growth restriction of the organism observed *in vitro* at 37°C in the presence of less than 2.5 mM Ca²⁺. Under such conditions the cells fail to synthesize bulk vegetative proteins although a series of stress proteins and virulence factors are expressed. The V antigen provides both active and passive immunity against experimental infection with F1⁺ strains [Lawton *et al.* (1963) *J. Immunol.* **91**: 179-184; Leary *et al.* (1995) *Infect. Immun.* **63**:
25 2854-2858; Nakajima *et al.* (1995) *Infect. Immun.* **63**: 3021-3029].

Vaccines which have focused on the capsule protein, F1, expressed either as the purified protein [Baker et al. (1952) *J. Immunol.* **68**: 131-145; Simpson et al. (1990) *Am. J. Trop. Med. Hyg.* **43**: 389-396] or in *aroA* mutant of *Salmonella typhimurium* [Oyston et al. (1995) *Infect. Immun.* **63**:563-568] protected mice against virulent *Y. pestis*.

- 5 However, *Y. pestis* strains lacking or deficient in F1 were isolated from immunized animals after challenge with F1⁺ organisms from wild rodents [Welkos et al. (1995) *Contrib. Microbiol. Immunol.* **13**:299-305] and from a fatal human case of plague [Winter et al. (1960) *Bull. WHO* **23**: 408-409]. Most importantly, F1⁻ strains are virulent in mice [Worsham et al. (1995) *Contrib. Microbiol. Immunol.* **13**: 325-327] and nonhuman
- 10 primates [Friedlander et al. (1995) *Clin. Infect. Dis.* **21**: (Suppl 2), S178-181] and we found in the present study that the human plague vaccine does not protect against infection with F1⁻ organisms. This implies that, as originally suggested by Burrows [Burrows, T. W. (1957) *Nature* **179**: 1246-1247], an improved plague vaccine protective against both F1⁻ and F1⁺ strains of *Y. pestis* must not rely solely on F1.
- 15 Therefore, there is a need to develop an improved vaccine protective against both F1⁻ and F1⁺ strains of *Y. pestis* suitable for human administration.

SUMMARY

- 20 The present invention is directed to a vaccine that satisfies this need. The vaccine of the present invention is protective against both F1⁻ and F1⁺ strains of *Y. pestis*. The vaccine of the present invention is composed of a fusion between a portion of the F1 protein and another protective immunogen, the V antigen. This invention is novel because it is a single constructed protein, F1-V, composed of two unique proteins, the entire F1
- 25 capsule antigen and V antigen. It induces an immunological response against both F1 protein and V antigen.

08699716 "082796

The invention was designed to be used in a vaccine affording protection against plague, and to solve the problem of protecting humans against both bubonic and pneumonic plague caused by infection by the subcutaneous (insect bite) and aerosol routes, respectively, with F1⁺ or F1⁻ plague organisms, or with strains which may vary in their V antigen.

The advantages of using this protein over the present whole cell vaccine are as follows:

- The current licensed vaccine does not protect mice against subcutaneous challenge with F1⁻ strains of *Y. pestis*, which have been shown to cause fatal disease in both humans and experimental animals infected by a peripheral, non-respiratory route. The new F1-V vaccine does protect mice against bubonic plague caused by subcutaneous challenge (insect bite) with F1⁻ organisms.

- The current licensed vaccine does not protect mice against pneumonic plague induced by aerosol challenge with F1⁻ strains of *Y. pestis*. The new F1-V vaccine does protect mice against pneumonic plague caused by aerosol challenge with F1⁻ strains.

- The current licensed vaccine does not protect mice against pneumonic plague when challenged by the respiratory route with F1⁺ strains of *Y. pestis*. The new F1-V vaccine does protect mice against pneumonic plague caused by aerosol challenge with F1⁺ strains.

- The new F1-V vaccine is expected to protect humans against pneumonic plague produced by strains of *Y. pestis*, either naturally occurring or genetically engineered, which may be altered in their content or composition of V antigen, but which still contain F1. This is because the F1-V vaccine also contains F1. The current licensed vaccine does not protect against pneumonic plague induced by either F1⁻ or F1⁺ organisms when given by the aerosol route.

- The new F1-V vaccine is composed of two antigens, both of which have been shown to be protective. The combination of both antigens should provide better protection

against F1⁺ strains than either F1 or V when used alone as vaccines. This is possible because the immunity induced by F1 and by V occur by different mechanisms which may be additive or synergistic.

5 - Approximately 8% of humans immunized with the current licensed human plague vaccine fail to develop an immune response to F1 [Marshall *et al.* (1974) *J. Inf. Dis.* 129:S26-S29]. These non-responders may well be at risk for development of plague. The inclusion of two different protective antigens in the same vaccine will help to eliminate the problem of non-responders to either of the individual antigens and so increase the overall efficacy of vaccination in a human population.

10 - The new F1-V vaccine is composed of highly purified recombinant proteins which are very well defined. This contrasts with the present human licensed vaccine composed of whole bacteria. The nature of the protective immunogen(s) in the present licensed vaccine is completely unknown. The present licensed vaccine is known to contain and induce antibodies to F1 but it does not induce antibodies to V antigen in mice, suggesting that V
15 antigen is absent.

 - The F1-V protein was constructed so that a single protein could be purified as a vaccine component rather than having to produce and purify F1 and V antigen separately. The purification of a single protein as opposed to two separate proteins could result in considerable savings when manufacturing a vaccine.

20 Therefore, it is an object of the present invention to provide a *Y. pestis* DNA fragment encoding 1563 bp of a fusion protein comprising the F1 protein fused at its carboxyl terminus to the amino terminus of the entire V antigen useful in the production of a diagnostic agent and a vaccine.

25 It is another object of the present invention to provide an amino acid sequence for *Y. pestis* F1-V protein encoding 521 amino acids.

It is another object of the invention to provide a recombinant vector comprising a vector and the above described DNA fragment for use as a DNA vaccine.

It is a further object of the present invention to provide a host cell transformed with any of the above-described recombinant DNA constructs for use as a live bacterial vaccine
5 when the host cell is a bacteria such as Salmonella, BCG, or a live viral vaccine when the host cell is a virus such as adenovirus, or Venezuelan Equine Encephalitis virus. These transformed cells, bacteria and viruses can also be used as a source for the *Y. pestis* F1-V protein.

10 It is another object of the present invention to provide a method for producing *Y. pestis* F1-V fusion protein which comprises culturing a host cell under conditions such that a recombinant vector comprising a vector and the *Y. pestis* F1-V protein DNA fragment is expressed and F1-V protein is thereby produced, and isolating F1-V protein for use as a vaccine or a diagnostic agent.

15

It is still another object of the invention to provide a purified *Y. pestis* F1-V protein useful as a vaccine and a diagnostic agent.

20 It is a further object of the present invention to provide an antibody to the above-described F1-V protein for use as a therapeutic agent and a diagnostic agent.

It is yet another object of the invention to provide a *Y. pestis* vaccine comprising a F1-V protein effective for eliciting an antigenic and immunogenic response resulting in the protection of a mammal against *Y. pestis* infection by subcutaneous and aerosol route.

25

It is yet another object of the present invention to provide a method for the diagnosis of *Y. pestis* infection comprising the steps of:

- (i) contacting a sample from an individual suspected of having the infection with antibodies which recognize F1-V protein; and
- (ii) detecting the presence or absence of a complex formed between *Y. pestis* F1 and/or V antigen and antibodies specific therefor.

It is a further object of the present invention to provide a diagnostic kit comprising a F1-V protein antibody and ancillary reagents suitable for use in detecting the presence *Y. pestis* in mammalian sputum, serum, or tissues.

It is yet another object of the present invention to provide a therapeutic method for the treatment or amelioration of symptoms of infection by *Y. pestis* and other species of *Yersinia* such as *Y. enterocolitica* and *Y. pseudotuberculosis*, said method comprising providing to an individual in need of such treatment an effective amount of sera from individuals immunized with F1-V protein in a pharmaceutically acceptable excipient.

It is further another object of the present invention to provide a therapeutic method for the treatment or amelioration of symptoms of *Yersinia* infection, said method comprising providing to an individual in need of such treatment an effective amount of antibodies against F1-V protein of *Yersinia pestis* and all or a portion of V antigen of *Yersinia enterocolitica* and *Yersinia pseudotuberculosis* in a pharmaceutically acceptable excipient.

It is still another object of the present invention to provide antigenic epitopes of F1-V protein, which are useful in peptide vaccine design.

BRIEF DESCRIPTION OF THE DRAWINGS

5 These and other features, aspects, and advantages of the present invention will become better understood with reference to the following description and appended claims, and accompanying drawings where:

 Figure 1 shows plasmid pF1V showing an in-frame insertion of the F1 and V open reading frames at the *NdeI* and *BamHI* sites of pET19b separated by an inframe *EcoRI* site.

10 F1-V was transcribed by using the T7 promoter (arrow) within pET19b after induction of T7 polymerase by isopropyl- β -D-thiogalactopyranoside (IPTG). The F1-V amino acid sequence begins with a His-Tag and enterokinase cleavage site derived from pET19b before the methionine start codon of F1. The F1 portion consists of 170 amino acids followed by two amino acids, asparagine and glutamine (encoded by the *EcoRI* site) and

15 the entire sequence of the V antigen. F1-V, therefore, has 521 amino acids with a predicted molecular mass of 57,926 daltons.

 Figure 2 shows the gel electrophoresis of F1-V fusion protein. The F1-V fusion protein expressed from pET19b was isolated with 6M urea as recommended (Novagen). Residual endotoxin was removed by passing F1-V over an endotoxin removing gel column

20 (Pierce, Rockford, IL). After the column, F1-V contained 202 endotoxin units per mg of protein by Limulus amoebocyte lysate assay (Sigma, St. Louis, MO) (1 endotoxin unit = 0.1 ng of *E. coli* 055:B5 LPS standard). Recombinant F1 and V were cloned separately and purified as described in Andrews *et al.* (1996) *Inf. Immun.* **64**:2180-2187. F1-V, V and F1 proteins (2 μ g each) were subjected to SDS-PAGE on 10% tricine gels (Novex, San

25 Diego, CA) and visualized by Coomassie brilliant blue staining.

DESCRIPTION

5 In one embodiment, the present invention relates to a DNA or cDNA segment which encodes *Y. pestis* recombinant F1-V protein consisting of the F1 protein fused at its carboxyl terminus to the amino terminus of the entire V antigen. The sequence of the 1563 nucleotide DNA segment is specified in SEQ ID NO: 1.

 DNA or polynucleotide sequences to which the invention also relates include
10 fragments of F1 or V containing protective epitopes [Motin *et al.* (1994) *Infect Immun.* 62:4192-4201].

 The derived polynucleotide is not necessarily physically derived from the nucleotide sequence shown in SEQ ID NO:1, but may be generated in any manner, including for example, chemical synthesis or DNA replication or reverse transcription or transcription,
15 which are based on the information provided by the sequence of bases in the region(s) from which the polynucleotide is derived. In addition, combinations of regions corresponding to that of the designated sequence may be modified in ways known in the art to be consistent with an intended use. The sequences of the present invention can be used in diagnostic assays such as hybridization assays and polymerase chain reaction assays for the detection
20 of F1 or V sequences of *Y. pestis*.

 In another embodiment, the present invention relates to a recombinant DNA molecule that includes a vector and a DNA sequence as described above. The vector can take the form of a plasmid such as pET19b, pMBac/pPBac, pSSV1 or any broad host
25 range expression vector such as viral vectors such as adenovirus or Venezuelan Equine Encephalitis virus and others known in the art.

In a further embodiment, the present invention relates to host cells stably transformed or transfected with the above-described recombinant DNA constructs. The host cell can be prokaryotic such as *Bacillus* or *E. coli*, or eukaryotic such as *Saccharomyces* or *Pichia*, or mammalian cells or insect cells. The vector containing the F1-V protein sequence is expressed in the bacteria and the expressed product used for diagnostic procedures or as a vaccine. Please see e.g., Maniatis, Fitch and Sambrook, Molecular Cloning: A Laboratory Manual (1982) or DNA Cloning, Volumes I and II (D. N. Glover ed. 1985) for general cloning methods. The DNA sequence can be present in the vector operably linked to a highly purified IgG molecule, an adjuvant, a carrier, or an agent for aid in purification of F1-V protein. The transformed or transfected host cells can be used as a source of DNA sequences described above. When the recombinant molecule takes the form of an expression system, the transformed or transfected cells can be used as a source of the protein described below.

In another embodiment, the present invention relates to a DNA sequence incorporated into a vector which can be used as a DNA vaccine in animals, including humans, or which can be used in a live bacterial or viral vaccine, e.g. *Salmonella*, BCG, adenovirus, or Venezuelan Equine Encephalitis virus.

In another embodiment, the present invention relates to a *Y. pestis* F1-V fusion protein having an amino acid sequence corresponding to SEQ ID NO: 2 and encompassing 521 amino acids or any allelic variation thereof.

A polypeptide or amino acid sequence derived from the amino acid sequence in SEQ ID NO:2, refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 2-5 amino acids, and more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence. The present invention relates to a F1-V protein where the F1 is fused by its carboxy-terminus to the amino-terminus of V antigen constructed

with a polyhistidine and enterokinase site and where the F1 signal sequence is present. Also embodied in this invention is a F1-V protein wherein the F1 is fused to the amino terminus of V, with or without any of the polyhistidine, enterokinase sites and the F1 signal sequence. In addition, a linker of additional amino acids can be fused in frame
5 between the F1 and V antigen (or V antigen and F1) sequences or fragments thereof for convenience as long as the changes do not affect the immunological activity of the fusion protein.

A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence, or the sequence in SEQ ID NO:1; it may be generated in
10 any manner, including for example, chemical synthesis, or expression of a recombinant expression system. In addition the polypeptide can be fused to other proteins or polypeptides for the purposes of transport or for secretion from the cell or for increasing the protective efficacy in a vaccine. Some examples include the *Yersinia* outer proteins (Yops) of *Yersinia* species or fragments thereof, or the amino-terminal protective antigen
15 binding domain of anthrax toxin lethal and edema factors to name a few.

In a further embodiment, the present invention relates to a method of producing F1-V protein which includes culturing the above-described host cells, under conditions such that the DNA fragment is expressed and F1-V is produced. F1-V can then be isolated using
20 methodology well known in the art or by the production method described below. F1-V protein can be used as a vaccine for immunity against infection with the *Y. pestis* or as a diagnostic tool for detection of *Y. pestis* infection. The transformed host cells can be used to analyze the effectiveness of drugs and agents which inhibit *Y. pestis*, such as host proteins or chemically derived agents or other proteins which may interact with the bacteria
25 to down-regulate or alter the expression of F1 protein or V antigen.

In another embodiment, the present invention relates to antibodies specific for the above-described F1-V protein. For instance, an antibody can be raised against the complete F1-V or against a portion thereof. Persons with ordinary skill in the art using standard methodology can raise monoclonal and polyclonal antibodies to F1-V of the present invention, or a unique portion thereof. Material and methods for producing antibodies are well known in the art (see for example Goding, in, Monoclonal Antibodies: Principles and Practice, Chapter 4, 1986). In addition, the protein or polypeptide can be fused to or combined with other proteins or polypeptides or adjuvants which increase its antigenicity, thereby producing higher titers of neutralizing antibody when used as a vaccine. Examples of such proteins or polypeptides include cholera toxin B subunit and any adjuvants or carriers safe for human use, such as aluminum hydroxide.

In a further embodiment, the present invention relates to a method of detecting the presence of *Y. pestis* infection or antibodies against *Y. pestis* in a sample. Using standard methodology well known in the art, a diagnostic assay can be constructed by coating on a surface (i.e. a solid support) for example, a microtitration plate or a membrane (e.g. nitrocellulose membrane), all or a unique portion of the F1-V protein described above, and contacting it with the serum of a person suspected of having plague. The presence of a resulting complex formed between F1-V protein and antibodies specific for either F1 or V in the serum can be detected by any of the known methods common in the art, such as fluorescent antibody spectroscopy or colorimetry. This method of detection can be used, for example, for the diagnosis of bubonic and pneumonic plague.

Similarly, antibodies to F1-V protein can be used in a rapid diagnostic assay to detect the presence of F1 and/or V antigen in the serum of patients infected with *Yersinia pestis*. Such a test may also be of value in the rapid diagnosis of infection in humans or animals with other *Yersinia* species by detection of V antigen in serum or tissue samples.

The ability of an individual to fight *Y. pestis* infection is dependent on the individual's ability to produce antibodies against *Y. pestis*. Diagnostic assays, similar to those described above, designed to measure the production of protective antibodies against F1-V can be used to measure an individual's response to receiving a plague vaccine.

5

In another embodiment, the present invention relates to a diagnostic kit which contains F1-V protein from *Y. pestis* and ancillary reagents that are suitable for use in detecting the presence of antibodies to *Y. pestis*, *Y. enterocolitica* and *Y.*

pseudotuberculosis in serum or a tissue sample. Tissue samples contemplated can be
10 rodents and human, or other mammals. Ancillary reagents would include standard anti-rodent or anti-human antibodies.

In another embodiment, the present invention relates to a vaccine for protection against *Y. pestis* infections by aerosol or subcutaneous route (insect bite). The vaccine comprises F1-V protein, or an immunogenic portion thereof, from a specific strain or
15 species of *Yersinia pestis*. It could also contain V antigen from *Y. enterocolitica* and *Y. pseudotuberculosis*. The vaccine can be prepared by inducing expression of a recombinant expression vector comprising F1-V protein sequence and purifying the resulting protein. The purified F1-V protein is prepared for administration to mammals by methods known in the art, which can include filtering to sterilize the solution, diluting the solution, adding an
20 adjuvant and stabilizing the solution. The vaccine can be lyophilized to produce a vaccine against *Y. pestis* in a dried form for ease in transportation and storage. Further, the vaccine may be prepared in the form of a mixed vaccine which contains the F1-V protein described above and at least one other antigen as long as the added antigen does not interfere with the effectiveness of the vaccine and the side effects and adverse reactions are
25 not increased additively or synergistically.

The vaccine may be stored in a sealed vial, ampule or the like. The present vaccine can generally be administered in the form of a liquid or suspension. In the case where the vaccine is in a dried form, the vaccine is dissolved or suspended in sterilized distilled water before administration. Generally, the vaccine may be administered orally, subcutaneously, 5 intradermally or intramuscularly but preferably intranasally in a dose effective for the production of neutralizing antibody and protection from infection or disease.

In another embodiment, the present invention relates to a method of reducing *Y. pestis* infection symptoms in a patient with bubonic or pneumonic plague by administering 10 to said patient an effective amount of F1-V protein antibodies including those made in humans, either polyclonal or combinations of monoclonals to F1 and V antigen, as described above. When providing a patient with F1-V antibodies, the dosage administered will vary depending upon such factors as the patient's age, weight, height, sex, general medical condition, previous medical history, etc. In general, it is desirable to provide the 15 recipient with a dosage of the above compounds which is in the range of from about 1 pg/kg to 500 mg/kg (body weight of patient), although a lower or higher dosage may be administered.

20 Described below are examples of the present invention which are provided only for illustrative purposes, and not to limit the scope of the present invention. In light of the present disclosure, numerous embodiments within the scope of the claims will be apparent to those of ordinary skill in the art.

25 The following Materials and Methods were used in the Examples below.

DNA methods. All restriction enzymes used were purchased from Life Technologies (Gaithersburg, Md.). Plasmid DNA samples were purified using a Qiagen

962280-9726980

plasmid purification kit (Qiagen, Inc., Chatsworth, Calif.) All oligonucleotide primers were synthesized on an Applied Biosystems model 391 DNA synthesizer (Foster City, Calif.) and the polymerase chain reaction (PCR) was performed using a DNA Thermal Cycler (Perkin-Elmer Cetus, Norwalk, Conn.). DNA sequencing was performed by the dideoxy-chain termination method [Sanger *et al.* (1977) *Proc. Natl. Acad. Sci.* **74**: 5463-5467] using [α -³⁵S]dATP (Amersham, Arlington Heights, Ill.). Genetic manipulations were performed by standard procedures [Sambrook *et al.* (1989) Molecular Cloning; A Laboratory Manual, 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.].

Analytical methods. Protein content was determined by the bicinchoninic acid-Lowry method with BSA as a standard. (Pierce). Purified F1-V was analyzed by sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis (PAGE) on 10% tricine gels (Novex, San Diego, Calif.) and visualized by Coomassie blue R-250 (Bio-Rad Laboratories, Hercules, Calif.) staining.

15

EXAMPLE 1

Construction, purification, and characterization of recombinant F1-V fusion protein.

To first isolate the F1 structural gene minus its stop codon, plasmid pYPR1 (kindly provided by T. Schwan, Rocky Mountain Laboratories, Hamilton, Mont.) containing the F1 operon [Simpson *et al.* (1990) *Am. J. Trop. Med. Hyg.* **43**:389-396] was used as template DNA in a PCR using as the forward oligonucleotide primer,

GGCGCGGCATATGAAAAAATCAGTTCC (SEQ ID NO:3), containing an internal

Nde I restriction site (underlined), and the reverse primer,

CTCGAATTCTTGGTTAGATACGGT (SEQ ID NO:4), containing an internal *Eco* R1 site

(underlined). The V antigen gene was then isolated by PCR using plasmid DNA from a

962280 9726980
pigmentation negative derivative (Pgm') of *Y. pestis* CO92, the forward oligonucleotide primer, CGCGAATTCATGATTAGAGCCTACGAA (SEQ ID NO: 5), containing an internal *Eco* RI site (underlined), and the reverse primer,

CGCGGATCCTCATTACCAGACGTGTCA (SEQ ID NO:6) containing an internal *Bam*

5 HI site (underlined). The purified F1 PCR product was then digested with *Nde* I and *Eco* RI while the V antigen purified PCR product was digested with *Eco* RI and *Bam* HI. Both restricted fragments were then ligated to the *Nde* I and *Bam* HI digested expression vector pET19b (Novagen, Madison, Wis.) and used to transform *Escherichia coli* strain BLR (Novagen) to create plasmid pF1V. The final protein contains an amino-terminal 10
10 histidines and enterokinase site from pET19b followed by the F1-V protein. The F1 portion consists of 170 amino acids followed by two amino acids, glutamic acid and phenylalanine (the *Eco* RI site) and the entire sequence of the V antigen. F1-V, therefore, has 521 amino acids with a predicted molecular mass of 57,926 daltons. The nucleotide
15 sequence of the F1-V portion of pF1V was verified by sequencing. Two nucleotide differences were found between the V sequence present in F1-V and that reported previously for the V antigen [Price *et al.* (1989) *J. Bacteriol.* 171: 5646-5653]. A G was replaced by an A at base 247 of the open reading frame, resulting in a change from an alanine to a threonine. At base 324, a G was replaced with a C with no change in the amino acid at that site.

20 The F1-V fusion protein expressed from pET19b was isolated with 6 M urea as recommended (Novagen). Residual endotoxin was removed by passing F1-V over an endotoxin removing gel column (Pierce, Rockford, Ill.). After the column, F1-V contained 202 endotoxin units per mg of protein by Limulus amoebocyte lysate assay (Sigma, St. Louis, Mo.) (1 endotoxin unit = 0.1 ng of *E. coli* 055:B5 LPS standard).

25 The recombinant F1-V fusion protein constructed in pET19b consists of the F1 protein fused at its carboxyl terminus to the amino terminus of the entire V antigen. F1-V had a relative molecular weight of 58,000 on SDS-PAGE (Fig. 2) which agreed with its

predicted size of 57,926. The F1-V protein reacted on Western blot with both rabbit polyclonal antibody to F1 as well as mouse monoclonal antibodies directed against V antigen (data not shown). Recombinant F1 and V antigen are also shown in Fig. 2.

- After establishing that F1-V had a relative molecular weight consistent with its DNA coding sequence and contained both F1 and V specific epitopes, we tested its ability to protect mice against plague.

EXAMPLE 2

- Animal immunization and challenge with *Y. pestis*.** Groups of female 8-10 week old Swiss Webster (Hsd:ND4) mice (Harlan Sprague Dawley, Indianapolis, Ind.) were immunized subcutaneously on days 0 and 28 with 0.2 ml of the F1-V, F1, or V antigen preparation adsorbed to the aluminum hydroxide adjuvant, Alhydrogel (1.3%, Superfos Biosector, Vedbaek, Denmark, 0.19 mg aluminum per dose), the human whole-cell plague vaccine U. S. Pharmacopeia (USP) (Greer Laboratories, Lenoir, NC) or Alhydrogel alone as a control. Serum obtained on day 58 after initial immunization was assayed for anti-F1 and anti-V IgG antibody by standard ELISA on individual animals and group geometric mean titers determined. Titers were determined as the reciprocal of the maximum dilution giving an absorbance greater than 0.1 units after subtraction of nonspecific binding in normal serum.
- The immunized animals were then challenged on day 78 by either the subcutaneous or aerosol route with wild-type F1⁺ *Y. pestis*, CO92 (kindly provided by T. Quan, Center for Disease Control, Ft. Collins, Colo.) or C12, an F1 isogenic derivative of CO92 with a deletion in the F1 structural gene [Worsham *et al.* (1995) *Contrib. Microbiol. Immunol.* 13: 325-327]. The inocula for s.c. and aerosol challenge were prepared and the animals challenged by s.c. and aerosol routes as previously described [Welkos *et al.* (1995) *Contrib. Microbiol. Immunol.* 13: 299-305]. The s.c. LD₅₀ is 9.1 and 1.9 CFU for F1⁺

C12 (37) and F1⁺ C092 (32) strains, respectively. The aerosol LD₅₀ is 1.1 x 10⁵ and 2 x 10⁴ CFU (32) for F1⁺ C12 and F1⁺ C092 strains, respectively.

All animal experiments were conducted in accordance with the *Guide for the Care and Use of Laboratory Animals* (Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council. 1985. Guidelines for the care and use of laboratory animals revised. National Institutes of Health publication no. 86-23. National Institutes of Health, Bethesda, MD) and animals were provided food and fresh water ad libitum during the experiment.

Efficacy of F1-V against bubonic plague.

In two separate experiments (Table 1), mice immunized with 13.6 µg of F1-V were protected (90-100% survival) against a subcutaneous challenge with a moderate (57 LD₅₀) or high (1.1 x 10⁶ LD₅₀) dose of the F1⁺ *Y. pestis* strain, C12, while all control animals died. Another group of animals immunized with 27.2 µg of F1-V all survived (100%) the high-dose challenge. As expected, F1 alone, when adsorbed to Alhydrogel, did not protect animals against challenge with the F1⁺ C12 strain. Animals given 10 µg of V were afforded the same degree of protection (90% survival) as with F1-V, against the high-dose challenge. Thus, the protective efficacy of the F1-V fusion protein against infection with an F1⁺ *Y. pestis* strain was equivalent at this challenge dose, to that provided by V alone.

In marked contrast, in a separate experiment, the current human, whole-cell plague vaccine USP, failed to protect against a low dose challenge; none of nine challenged animals survived.

TABLE 1 Efficacy of F1-V vaccination against a lethal subcutaneous *Y. pestis* infection of mice.

	Treatment Group	Strain	Challenge dose ^a	Survivors/total
5	Alhydrogel alone	F1 ⁻ C12	57	0/10
	13.6 µg F1-V	"	57	10/10
	10 µg F1	"	60	0/10
	Plague USP ^b	"	124	0/9
	Alhydrogel alone	"	1.1x10 ⁶	0/10
10	13.6 µg F1-V	"	1.1x10 ⁶	9/10
	27.2 µg F1-V	"	1.1x10 ⁶	10/10
	10 µg V	"	1.1x10 ⁶	9/10

^aNumber of LD₅₀

15 ^bThe licensed, human, whole-cell plague vaccine.

Efficacy of F1-V against pneumonic plague.

We next determined the efficacy of F1-V against pneumonic plague induced by an aerosol challenge (Table 2). Mice immunized with 13.6 µg of F1-V were completely
 20 protected (100% survival) against a moderate (91 LD₅₀) or high (590 LD₅₀) aerosol dose of the F1⁻ *Y. pestis* strain, C12. Animals given 27.2 µg of F1-V were also completely protected against the high-dose challenge. Similarly, V protected animals exposed to the high-dose aerosol challenge, with 80% of animals surviving. However, as with the s.c. challenge, the plague vaccine USP failed to protect against fatal pneumonic plague; none of
 25 eight challenged animals survived. Thus, the whole-cell plague vaccine USP failed to protect mice against challenge with the F1⁻ strain by either the s.c. or aerosol route, while it does protect against s.c. challenge [Andrews *et al.* (1996) *Infect. Immun.* 64:2180-2187;

- Simpson *et al.* (1990) *Am. J. Trop. Med. Hyg.* **43**:389-396] and partially protects with prolongation of time to death against aerosol challenge with F1⁺ strains [Andrews *et al.*, *ibid.*; Pitt *et al.* (1994) Abstr. E-45. In: Abstracts of the 94th General Meeting of the American Society for Microbiology 1994. American Society for Microbiology, Washington, D.C.; Smith and Packman (1966) *Brit. J. Exp. Path.* **47**: 25-34]. This differential protection against F1⁺ and F1⁻ strains and the absence of an immune response to V antigen in the plague vaccine USP group (Table 2) in agreement with other studies [Chen *et al.* (1961) *J. Immunol.* **87**:64-71; Williamson *et al.* (1995) *FEMS Immunol. Med. Microbiol.* **12**:223-230], strongly suggest that the major protective immunogen in the plague vaccine USP is F1 and that V antigen and other possible immunogens are absent.
- The F1-V also protected (100% survival) against a high-dose aerosol challenge with the F1⁺ *Y. pestis*, CO92 strain isolated from a fatal human pneumonic case. While these studies were in progress, a report described the increased effectiveness of co-immunization with F1 and V antigen in protecting against subcutaneous challenge with an F1⁺ strain [Williamson *et al.*, *ibid.*]. No studies were performed with F1⁻ strains or against an aerosol challenge.

TABLE 2. Efficacy of F1-V vaccination against a lethal aerosol *Y. pestis* infection of mice

Treatment Group	Strain	Challenge dose ^a	Survivors/Total	Geometric mean antibody titer ^b	
				F1	V
Alhydrogel alone	F1 ⁻ C12	91	0/9	NT ^c	NT
13.6 µg F1-V	"	91	10/10	NT	NT
Alhydrogel alone	"	590	0/14	<640	<640

TABLE 2 Continued. Efficacy of F1-V vaccination against a lethal aerosol *Y. pestis* infection of mice

5	Treatment Group	Strain	Challenge dose ^a	Survivors/Total	Geometric mean antibody titer ^b	
					F1	V
10	13.6 µg F1-V	"	590	10/10	66,540	432,376
	27.2 µg F1-V	"	590	10/10	108,094	432,376
	10 µg V	"	590	8/10	NT	655,360
	Plague USP ^d	"	590	0/8	55,738	<640
	Alhydrogel alone	F1 ⁺ CO92	761	1/10	NT	NT
	13.6 µg F1-V	"	761	10/10	NT	NT

15 ^aNumber of LD₅₀

^bSerum obtained on day 58 after the initial vaccine dose was assayed for anti-F1 and anti-V IgG antibody by ELISA on individual animals and group geometric mean titers determined as described in Materials and Methods.

^cNot tested.

20 ^dThe licensed, human, whole-cell plague vaccine.

Our results clearly demonstrate that mice immunized with a fusion protein, consisting of the F1 capsular antigen fused at its carboxyl terminus to the amino terminus of the V antigen of *Y. pestis*, were provided with excellent protection against both parenteral and aerosol challenge with an F1⁻ *Y. pestis* strain. The F1-V fusion protein also protected mice against an aerosol challenge with an F1⁺ *Y. pestis* strain. Other workers, as noted previously, showed that V antigen protects animals against challenge with virulent

F1⁺ strains, results which we have confirmed and extended to aerosol challenge. This raises the issue of whether the F1 portion of F1-V is immunogenic and contributes to the protection against challenge with the F1⁺ *Y. pestis* strain CO92. Several lines of evidence suggest that the F1 portion is immunogenic and protective. The development of high levels of antibody to F1 after immunization with F1-V (Table 2) and the numerous reported studies showing a strong correlation between the level of antibody to F1 and protection against infection [Williams and Cavanaugh (1979) Bull. WHO 57:309-313] suggest the F1 portion of F1-V helps protect against challenge with F1⁺ *Y. pestis*. Further support for a protective role for the F1 portion of F1-V was provided by our observations with a smaller F1-V fusion protein we constructed that consisted of the entire F1 gene fused at its carboxyl terminus to amino acids 168 to 275 of V antigen. This fusion protein was unable to effectively immunize mice against a subcutaneous challenge with the F1⁺ *Y. pestis* C12 strain (3 survivors out of 10 mice challenged with 55 LD₅₀, data not shown), showing that the V segment of the fusion protein was not protective. However, this same fusion protein was able to protect mice against the F1⁺ *Y. pestis* CO92 strain (10 survivors out of 10 mice challenged subcutaneously with 63 LD₅₀, data not shown). These results indicate the F1 portion of this fusion protein was immunogenic, enabling mice to survive challenge against infection with the F1⁺ *Y. pestis* CO92, while the V segment failed to elicit a significant protective immune response against challenge with the V antigen expressing but F1⁻ *Y. pestis* C12 strain. The value of a combined immune response to both F1 and V for combating infection with F1⁺ *Y. pestis* strains is supported by the studies of Burrows and Bacon [Burrows, T. W. (1963) *Ergeb. Mikrobiol. Immunitätsforsch. Exp. Ther.* 37:59-113; Burrows and Bacon (1958) *Brit. J. Exp. Pathol.* 39: 278-291] who found that serum from rabbits immunized with attenuated strains of *Y. pestis* expressing both F1 and V antigen provided better passive protection in mice against wild type F1⁺ *Y. pestis* than serum from rabbits immunized with attenuated *Y. pestis* strains expressing only V antigen. It is also supported by the recent study showing increased protection by co-immunization

of F1 with V antigen [Williamson *et al.*, *ibid.*]. Better protection by antibody directed against both F1 and V might occur by counteracting both the anti-phagocytic activity associated with F1 [Burrows, T. W., *ibid.*] and the virulence-enhancing activity associated with secreted V antigen [Nakajima *et al.* (1995) *Infect. Immun.* **63**: 3021-3029].

5 An additional advantage of a vaccine containing both F1 and V is that F1 should protect against variant strains of *Y. pestis* which might be altered in the amount or composition of V antigen, in a manner analogous to that by which V protects against F1⁻ strains. Indeed, isolates deficient in V, determined immunologically, have been cultured from immunized animals infected with V-containing, wild-type *Y. pestis* [Williams *et al.* 10 (1974) *Trans. Roy. Soc. Trop. Med. Hyg.* **68**:171]. Furthermore, variability in the structural gene for V has been described for *Y. pseudotuberculosis* [Motin *et al.* (1994) *Infect. Immun.* **62**:4192-4201], although to date, this has not been reported for *Y. pestis*.

Another potential advantage of an F1-V multicomponent vaccine would be to protect individuals who may be non-responders to one component of a vaccine. Indeed, 15 some recipients of the current plague vaccine USP fail to develop an antibody response to F1 [Marshall *et al.* (1974) *J. Infect. Dis.* **129**:S26-S29]. A similar approach to the development of recombinant multicomponent vaccines may be of value against other infectious diseases.

In summary, we constructed an F1-V fusion protein to provide optimal protective 20 immunity against pneumonic as well as bubonic plague due to either wild-type F1⁺ *Y. pestis* or fully virulent F1⁻ *Y. pestis* strains which may occur naturally or may develop after infection of vaccinated individuals. The new vaccine was shown to be effective against both F1⁻ as well as F1⁺ *Y. pestis* strains. Most importantly, it prevented both fatal pneumonic as well as bubonic infection. This vaccine candidate may lead to the 25 development of an improved human plague vaccine.

SEQUENCE LISTING

- 5 (1) GENERAL INFORMATION:
(i) APPLICANT: David G. Heath
Arthur M. Friedlander
George W. Anderson
Susan L. Welkos
- 10 (ii) TITLE OF INVENTION: Recombinant F1-V Plague Vaccine
- (iii) NUMBER OF SEQUENCES: 6
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: John Moran
(B) STREET: USA MPMC - MPMC-JA
(C) CITY: FORT DETRICK, FREDERICK
(D) STATE: MARYLAND
(E) COUNTRY: USA
20 (F) ZIP: 21702-5012
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: Apple Macintosh
25 (C) OPERATING SYSTEM: Macintosh 7.5
(D) SOFTWARE: Microsoft Word 6.0
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
30 (B) FILING DATE:
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER:
35 (B) FILING DATE:
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Moran, John
(B) REGISTRATION NUMBER: 26,313
40 (C) REFERENCE/DOCKET NUMBER:
- (ix) TELECOMMUNICATION INFORMATION
(A) TELEPHONE: (301) 619-2065
(B) TELEFAX: (301) 619-7714
45
- (2) INFORMATION FOR SEQ ID NO: 1:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1566 bp
50 (B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:1:

5	ATGGGCCATC	ATCATCATCA	TCATCATCAT	CATCACAGCA	40
	GCGGCCATAT	CGACGACGAC	GACAAGCATA	TGAAAAAAT	80
	CAGTTCCGTT	ATCGCCATTG	CATTATTTGG	AACTATTGCA	120
10	ACTGCTAATG	CGGCAGATTT	AACTGCAAGC	ACCACTGCAA	160
	GCGCAACTCT	TGTTGAACCA	GCCCGCATCA	CTCTTACATA	200
15	TAAGGAAGGC	GCTCCAATTA	CAATTATGGA	CAATGGAAAC	240
	ATCGATACAG	AATTACTTGT	TGGTACGCTT	ACTCTTGGCG	280
	GCTATAAAAC	AGGAACCACT	AGCACATCTG	TTAACTTTAC	320
20	AGATGCCGCG	GGTGATCCCA	TGTACTTAAC	ATTTACTTCT	360
	CAGGATGGAA	ATAACCACCA	ATTCACTACA	AAAGTGATTG	400
25	GCAAGGATTC	TAGAGATTTT	GATATCTCTC	CTAAGGTAAA	440
	CGGTGAGAAC	CTTGTGGGGG	ATGACGTCGT	CTTGGCTACG	480
	GGCAGCCAGG	ATTTCTTTGT	TCGCTCAATT	GGTTCCAAAG	520
30	GCGGTAAACT	TGCAGCAGGT	AAATACACTG	ATGCTGTAAC	560
	CGTAACCGTA	TCTAACCAAG	AATTCATGAT	TAGAGCCTAC	600
35	GAACAAAACC	CACAACATTT	TATTGAGGAT	CTAGAAAAAG	640
	TTAGGGTGGA	ACAACCTACT	GGTCATGGTT	CTTCAGTTTT	680
	AGAAGAATTG	GTTTCAGTTAG	TCAAAGATAA	AAATATAGAT	720
40	ATTTCCATTA	AATATGATCC	CAGAAAAGAT	TCGGAGGTTT	760
	TTGCCAATAG	AGTAATTACT	GATGATATCG	AATTGCTCAA	800
45	GAAAATCCTA	GCTTATTTTC	TACCCGAGGA	TACCATTCTT	840
	AAAGGCGGTC	ATTATGACAA	CCAACTGCAA	AATGGCATCA	880
	AGCGAGTAAA	AGAGTTCCTT	GAATCATCGC	CGAATACACA	920
50	ATGGGAATTG	CGGGCGTTCA	TGGCAGTAAT	GCATTTCTCT	960

	TTAACCGCCG	ATCGTATCGA	TGATGATATT	TTGAAAGTGA	1000
	TTGTTGATTC	AATGAATCAT	CATGGTGATG	CCCGTAGCAA	1040
5	GTTGCGTGAA	GAATTAGCTG	AGCTTACCGC	CGAATTAAAG	1080
	ATTTATTCAG	TTATTCAAGC	CGAAATTAAT	AAGCATCTGT	1120
	CTAGTAGTGG	CACCATAAAAT	ATCCATGATA	AATCCATTAA	1160
10	TCTCATGGAT	AAAAATTTAT	ATGGTTATAC	AGATGAAGAG	1200
	ATTTTTAAAG	CCAGCGCAGA	GTACAAAATT	CTCGAGAAAA	1240
15	TGCCTCAAAC	CACCATTCAG	GTGGATGGGA	GCGAGAAAAA	1280
	AATAGTCTCG	ATAAAGGACT	TTCTTGGGAG	TGAGAATAAA	1320
	AGAACCGGGG	CGTTGGGTAA	TCTGAAAAAC	TCATACTCTT	1360
20	ATAATAAAGA	TAATAATGAA	TTATCTCACT	TTGCCACCAC	1400
	CTGCTCGGAT	AAGTCCAGGC	CGCTCAACGA	CTTGGTTAGC	1440
25	CAAAAAACAA	CTCAGCTGTC	TGATATTACA	TCACGTTTTA	1480
	ATTCAGCTAT	TGAAGCACTG	AACCGTTTCA	TTCAGAAATA	1520
	TGATTCACTG	ATGCAACGTC	TGCTAGATGA	CACGTCTGGT	1560
30	AAATGA				1566

(3) INFORMATION FOR SEQ ID NO:2:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 521
 (B) TYPE: Amino acid
 (C) STRANDEDNESS: Single
 40 (D) TOPOLOGY: Linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:2:

45	Met	Gly	His	His	His	His	His	His	His	His	His	His	Ser	Ser	Gly
	1				5						10				15
	His	Ile	Asp	Asp	Asp	Asp	Lys	His	Met	Lys	Lys	Ile	Ser	Ser	Val
					20					25					30
50	Ile	Ala	Ile	Ala	Leu	Phe	Gly	Thr	Ile	Ala	Thr	Ala	Asn	Ala	Ala

	35	40	45
	Asp Leu Thr Ala Ser Thr Thr Ala Thr	Ala Thr Leu Val Glu Pro	
	50	55	60
5	Ala Arg Ile Thr Leu Thr Tyr Lys Glu	Gly Ala Pro Ile Thr Ile	
	65	70	75
10	Met Asp Asn Gly Asn Ile Asp Thr Glu	Leu Leu Val Gly Thr Leu	
	80	85	90
	Thr Leu Gly Gly Tyr Lys Thr Gly Thr	Thr Ser Thr Ser Val Asn	
	95	100	105
15	Phe Thr Asp Ala Ala Gly Asp Pro Met	Tyr Leu Thr Phe Thr Ser	
	110	115	120
	Gln Asp Gly Asn Asn His Gln Phe Thr	Thr Lys Val Ile Gly Lys	
	125	130	135
20	Asp Ser Arg Asp Phe Asp Ile Ser Pro	Lys Val Asn Gly Glu Asn	
	140	145	150
25	Leu Val Gly Asp Asp Val Val Leu Ala	Thr Gly Ser Gln Asp Phe	
	155	160	165
	Phe Val Arg Ser Ile Gly Ser Lys Gly	Gly Lys Leu Ala Ala Gly	
	170	175	180
30	Lys Tyr Thr Asp Ala Val Thr Val Thr	Val Ser Asn Gln Glu Phe	
	185	190	195
	Met Ile Arg Ala Tyr Glu Gln Asn Pro	Gln His Phe Ile Glu Asp	
	200	205	210
35	Leu Glu Lys Val Arg Val Glu Gln Leu	Thr Gly His Gly Ser Ser	
	215	220	225
40	Val Leu Glu Glu Leu Val Gln Leu Val	Lys Asp Lys Asn Ile Asp	
	230	235	240
	Ile Ser Ile Lys Tyr Asp Pro Arg Lys	Asp Ser Glu Val Phe Ala	
	245	250	255
45	Asn Arg Val Ile Thr Asp Asp Ile Glu	Leu Leu Lys Lys Ile Leu	
	260	265	270
	Ala Tyr Phe Leu Pro Glu Asp Thr Ile	Leu Lys Gly Gly His Tyr	
	275	280	285
50	Asp Asn Gln Leu Gln Asn Gly Ile Lys	Arg Val Lys Glu Phe Leu	
	290	295	300
	Glu Ser Ser Pro Asn Thr Gln Trp Glu	Leu Arg Ala Phe Met Ala	

		305		310		315
	Val Met His Phe	Ser Leu Thr Ala Asp Arg	Ile Asp Asp Asp	Ile		
		320		325		330
5	Leu Lys Val Ile	Val Asp Ser Met Asn His	His Gly Asp Ala Arg			
		335		340		345
10	Ser Lys Leu Arg	Glu Glu Leu Ala Glu	Leu Thr Ala Glu Leu Lys			
		350		355		360
	Ile Tyr Ser Val	Ile Gln Ala Glu Ile	Asn Lys His Leu Ser Ser			
		365		370		375
15	Ser Gly Thr Ile	Asn Ile His Asp Lys	Ser Ile Asn Leu Met Asp			
		380		385		390
	Lys Asn Leu Tyr	Gly Tyr Thr Asp Glu	Glu Ile Phe Lys Ala Ser			
		395		400		405
20	Ala Glu Tyr Lys	Ile Leu Glu Lys Met	Pro Gln Thr Thr Ile Gln			
		410		415		420
25	Val Asp Gly Ser	Glu Lys Lys Ile Val	Ser Ile Lys Asp Phe Leu			
		425		430		435
	Gly Ser Glu Asn	Lys Arg Thr Gly Ala	Leu Gly Asn Leu Lys Asn			
		440		445		450
30	Ser Tyr Ser Tyr	Asn Lys Asp Asn Asn	Glu Leu Ser His Phe Ala			
		455		460		465
	Thr Thr Cys Ser	Asp Lys Ser Arg Pro	Leu Asn Asp Leu Val Ser			
		470		475		480
35	Gln Lys Thr Thr	Gln Leu Ser Asp Ile	Thr Ser Arg Phe Asn Ser			
		485		490		495
40	Ala Ile Glu Ala	Leu Asn Arg Phe Ile	Gln Lys Tyr Asp Ser Val			
		500		505		510
	Met Gln Arg Leu	Leu Asp Asp Thr Ser	Gly Lys			
		515		520		

45

(4) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

50

- (A) LENGTH: 28 bp
- (B) TYPE: Nucleic acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGCGCGGCAT ATGAAAAAAA TCAGTTCC 28

5

(5) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 24 bp
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:4:

15

CTCGAATTCT TGGTTAGATA CGGT 24

(6) INFORMATION FOR SEQ ID NO:5:

20

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 27 bp
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:5:

30

CGCGAATTCA TGATTAGAGC CTACGAA 27

(7) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 28 bp
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear

(ii) SEQUENCE DESCRIPTION: SEQ ID NO:6:

40

CGCGGATCCT CATTTACCAG ACGTGTCA 28

45

50

What is claimed is:

1. An isolated and purified F1-V DNA fragment which encodes all or a
5 portion of F1 capsular antigen of *Yersinia pestis* and all or a portion of V antigen of
Yersinia pestis.
2. An isolated and purified DNA fragment according to claim 1 wherein, F1
capsular antigen of *Y. pestis* is fused at its carboxyl terminal to amino acid terminus of V
10 antigen of *Yersinia pestis*.
3. An isolated and purified DNA fragment according to claim 2, wherein
said fragment has the sequence of SEQ ID NO: 1 or a portion thereof, or an allelic portion
thereof.
15
4. An isolated and purified F1-V DNA fragment according to claim 3 which
encodes 521 amino acids or a portion thereof.
5. The F1-V DNA fragment according to claim 4, wherein said DNA
20 fragment encodes the amino acid sequence according to SEQ ID NO: 2 or a portion thereof.
6. A F1-V DNA fragment according to claim 2, comprising at least 30
nucleotides of the sequence set forth therein.
- 25 7. A recombinant DNA construct comprising:
(i) a vector, and

(ii) an isolated and purified F1-V DNA fragment which encodes all or a portion of F1 capsular antigen of *Yersinia pestis* and all or a portion of V antigen of *Yersinia pestis*.

5 8. A recombinant DNA construct according to claim 7, wherein said DNA fragment encodes the amino acid sequence specified in SEQ ID NO:2.

9. A recombinant DNA construct according to claim 7, wherein said vector is an expression vector.

10 10. The recombinant DNA construct according to claim 7, wherein said vector is a prokaryotic vector.

15 11. A recombinant DNA construct according to claim 8 wherein said construct is pF1V.

12. A recombinant DNA construct according to claim 7, wherein said vector is a eukaryotic vector.

20 13. A host cell transformed with a recombinant DNA construct comprising:
 (i) a vector, and
 (ii) a F1-V DNA fragment which encodes 521 amino acids of
 F1-V which encodes all or a portion of F1 capsular antigen of
 Yersinia pestis and all or a portion of V antigen of *Yersinia pestis*.

25 14. A host cell according to claim 13, wherein said cell is prokaryotic.

15. A host cell according to claim 14, wherein said host cell is *E. coli BLR*.

16. A host cell according to claim 13, wherein said host is a virus.

5 17. A method for producing F1-V protein which comprises culturing host cells transformed with an expression vector containing a DNA fragment encoding all or a portion of F1 capsular protein of *Yersinia pestis* and all or a portion of V antigen of *Yersinia pestis*, under conditions such that said DNA fragment is expressed and said F1-V protein is thereby produced, and isolating said F1-V protein.

10 18. A F1-V protein comprising all or a portion of F1 capsular antigen of *Yersinia pestis* and all or a portion of V antigen of *Yersinia pestis*.

15 19. A F1-V protein according to claim 18 wherein said F1 capsular protein is fused at its carboxyl terminus to amino acid terminus of V antigen and having the amino acid sequence specified in SEQ ID NO:2.

20 20. An antibody to a peptide having the amino acid sequence specified in SEQ ID NO:2 or a portion thereof.

25 21. A vaccine protective against naturally occurring or genetically engineered *Yersinia pestis* subcutaneous and aerosol infection comprising F1-V protein comprising all or a portion of F1 capsular antigen of *Yersinia pestis* and all or a portion of V antigen of *Yersinia pestis* capable of eliciting protective antibodies against *Y. pestis* infection in a pharmaceutically acceptable excipient in a pharmaceutically acceptable amount.

22. A vaccine protective against naturally occurring or genetically engineered *Yersinia pestis* according to claim 21 wherein, said *Y. pestis* is F1 deficient and V antigen altered.

5 23. A vaccine protective against naturally occurring or genetically engineered *Yersinia pestis* according to claim 21 wherein, said *Y. pestis* is F1 expressing and V antigen altered.

24. A vaccine for *Yersinia* infections comprising a F1-V protein comprising
10 all or a portion of F1 capsular antigen of *Yersinia pestis* and all or a portion of V antigen of *Yersinia pestis* and all or a portion of *Yersinia enterocolitica* V antigen and all or a portion of *Yersinia pseudotuberculosis* V antigen capable of eliciting protective antibodies against *Yersinia* infection in a pharmaceutically acceptable excipient in a pharmaceutically acceptable amount.

15 25. A *Y. pestis* infection diagnostic kit comprising an antibody to a peptide having the amino acid sequence specified in SEQ ID NO:2 or a portion thereof and ancillary reagents suitable for use in detecting the presence or absence of F1 or V in a mammalian sample.

20 26. A therapeutic method for the treatment or amelioration of a *Y. pestis* infection comprising administering to an individual having such an infection an effective amount of antibodies to a peptide having the amino acid sequence specified in SEQ ID NO:2 or a portion thereof in a pharmaceutically acceptable dose in a pharmaceutically
25 acceptable excipient.

27. A method for the diagnosis of *Y. pestis* infections comprising the steps
of:

(i) contacting a sample from an individual suspected of having a *Y. pestis*
infection with an antibody to a F1-V protein comprising all or a portion of F1 capsular
5 antigen of *Yersinia pestis* and all or a portion of V antigen of *Yersinia pestis*.; and

(ii) detecting the presence or absence of *Y. pestis* infection by detecting the
presence or absence of a complex formed between said antibody and F1 and V antigens
present in said sample.

10 28. A method for the diagnosis of *Y. pestis* infection comprising the steps
of :

(i) contacting a sample from an individual suspected of having a *Y. pestis*
infection with a F1-V protein comprising all or a portion of F1 capsular antigen of *Yersinia*
pestis and all or a portion of V antigen of *Yersinia pestis*.; and

15 (ii) detecting the presence or absence of a *Y. pestis* infection by detecting the
presence or absence of a complex formed between the F1-V protein and antibodies specific
for F1 or V in the sample.


20

25

ABSTRACT

- 5 A recombinant vaccine is described which is composed of one recombinant protein comprising F1 and V antigen of *Yersinia pestis*. This vaccine protects experimental mice against pneumonic as well as bubonic plague produced by either F1⁺ or F1⁻ strains of *Y. pestis* and may provide the basis for an improved human plague vaccine.


693716



pF1V

08699716-086996

1699716



03699716-082795